

AD-A152 856

A BILATERALLY DETERMINISTIC RHO-MIXING STATIONARY
RANDOM SEQUENCE(U) NORTH CAROLINA UNIV AT CHAPEL HILL
CENTER FOR STOCHASTIC PROCESSES R C BRADLEY FEB 85

1/1

UNCLASSIFIED

TR-91 AFOSR-TR-85-0348 F49620-82-C-0009


F/G 12/1

NL

END

FILMED


DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

CENTER FOR STOCHASTIC PROCESSES**AD-A152 856**

Department of Statistics
University of North Carolina
Chapel Hill, North Carolina



A BILATERALLY DETERMINISTIC ρ -MIXING
STATIONARY RANDOM SEQUENCE

by

Richard C. Bradley

Technical Report No. 91

February 1985

DTIC
ELECTE
APR 26 1985
S **B** **D**

DTIC FILE COPY

A BILATERALLY DETERMINISTIC ρ -MIXING
STATIONARY RANDOM SEQUENCE

by

Richard C. Bradley

Department of Mathematics
Indiana University
Bloomington
Indiana 47405

and

Center for Stochastic Processes
Department of Statistics
University of North Carolina
Chapel Hill, North Carolina 27514

Abstract

→ A (non-degenerate) strictly stationary sequence $(X_k, k \in \mathbb{Z})$ of random variables is constructed such that the ρ -mixing (maximal correlation mixing) condition is satisfied and each X_{sub-k} is measurable with respect to the double tail σ -field.

sigma

K

DTIC
ELECTE
S APR 26 1985 D
B

- A -

Parts of this work were done while the author was a visitor at the Mathematical Centre in Amsterdam and the Center for Stochastic Processes in Chapel Hill.

Research partially supported by AFOSR grant #F49620 82 C 0009 and by NSF grant DMS 84-01021.

DISTRIBUTION STATEMENT A

Approved for public release
Distribution Unlimited

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited.		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) 91			5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR- 85 - 0348		
6a. NAME OF PERFORMING ORGANIZATION University of North Carolina		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION Air Force Office of Scientific Research		
6c. ADDRESS (City, State and ZIP Code) Statistics Department, Phillips Hall 039-A Chapel Hill NC 27514		7b. ADDRESS (City, State and ZIP Code) Directorate of Mathematical & Information Sciences, Bolling AFB DC 20332-6448			
8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR		8b. OFFICE SYMBOL (If applicable) NM	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F49620-82-C-0009		
8c. ADDRESS (City, State and ZIP Code) Bolling AFB DC 20332-6448		10. SOURCE OF FUNDING NOS.			
		PROGRAM ELEMENT NO. 61102F	PROJECT NO. 2304	TASK NO. A5	WORK UNIT NO.
11. TITLE (Include Security Classification) A BILATERALLY DETERMINISTIC p-MIXING STATIONARY RANDOM SEQUENCE					
12. PERSONAL AUTHOR(S) Richard C. Bradley					
13a. TYPE OF REPORT Technical		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Yr., Mo., Day) FEB 85	
15. PAGE COUNT 15					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB. GR.			
19. ABSTRACT (Continue on reverse if necessary and identify by block number) A (non-degenerate) strictly stationary sequence $(X_k, k \in \mathbb{Z})$ of random variables is constructed such that the ρ -mixing (maximal correlation mixing) condition is satisfied and each X_k is measurable with respect to the double tail σ -field.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL MAJ Brian W. Woodruff			22b. TELEPHONE NUMBER (Include Area Code) (202) 767- 5027		22c. OFFICE SYMBOL NM

1. Introduction

Suppose $(X_k, k \in \mathbb{Z})$ is a strictly stationary sequence of random variables on a probability space (Ω, \mathcal{F}, P) . For $-\infty \leq J \leq L \leq \infty$ let \mathcal{F}_J^L denote the σ -field of events generated by $(X_k, J \leq k \leq L)$. For the sequence (X_k) the "past" and "future" tail σ -fields are respectively $\bigcap_{n=1}^{\infty} \mathcal{F}_{-\infty}^{-n}$ and $\bigcap_{n=1}^{\infty} \mathcal{F}_n^{\infty}$, and the "double" tail σ -field is $\bigcap_{n=1}^{\infty} (\mathcal{F}_{-\infty}^{-n} \vee \mathcal{F}_n^{\infty})$. The sequence (X_k) is said to be "bilaterally deterministic" if each of the r.v.'s X_k is measurable with respect to the double tail σ -field.

Several authors have discussed examples of strictly stationary sequences (X_k) which have a non-trivial double tail σ -field - or are even bilaterally deterministic - such that the past and future tail σ -fields are each trivial. Olshen [5] gives an example with the X_k 's in essence being real-valued. In Gurevič [2] and in Ornstein and Weiss [6], classes of finite-state examples are constructed, some of these examples being isomorphic to a Bernoulli shift. Also [6] refers to similar work of Furstenberg. Here we shall examine the question of what strong mixing conditions can be satisfied by bilaterally deterministic stationary sequences of real-valued r.v.'s. This work arose from a question posed to the author by H. Berbee as to whether the (Rosenblatt) strong mixing condition implied a trivial double tail σ -field.

For any two σ -fields A and B define the "maximal correlation":

$$\rho(A, B) := \sup_{f \in L^2(A), g \in L^2(B)} |\text{Corr}(f, g)|$$

(In the notation $L^2(\cdot)$ here, only real-valued random variables are included.) For a given strictly stationary sequence $(X_k, k \in \mathbb{Z})$, define for each $n \in \mathbb{N}$,

$\rho(n) := \rho(F_{-\infty}^0, F_n^\infty)$. The sequence (X_k) is said to be " ρ -mixing" if $\rho(n) \rightarrow 0$ as $n \rightarrow \infty$. The following theorem will be proved:

Theorem 1: Suppose c_1, c_2, c_3, \dots is a sequence of positive numbers. Then there exists a bilaterally deterministic strictly stationary sequence $(X_k, k \in \mathbb{Z})$ of non-degenerate real-valued random variables such that $\forall n \in \mathbb{N}, \rho(n) \leq c_n$.

In particular, a (non-degenerate) bilaterally deterministic strictly stationary sequence (X_k) can be ρ -mixing with an arbitrarily fast rate of convergence of $\rho(n)$ to 0. Now ρ -mixing implies the (Rosenblatt) strong mixing condition, which in turn implies that the past and future tail σ -fields are each trivial. (See e.g. [4, pp. 301-306, Def. 17.1.1, Theorem 17.1.1, and Def. 17.2.1].) Among the mixing conditions in the literature which fail to imply a trivial double tail σ -field, there does not seem to be any that is stronger than ρ -mixing. The ρ -mixing condition plays a natural role in central limit theory for dependent r.v.'s; see e.g. Ibragimov [3].

Theorem 1 is proved in Section 3. In Section 2 some preliminary work is done in preparation for that proof. Our construction in Sections 2 and 3 is similar to (but more complicated than) Olshen's; his construction (see [5, p. 155, lines 5-10, and p. 156, lines 4-12]) contains some key ideas used in our construction.



Accession For	
NTIS	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

2. Preliminaries

The following result of Csaki and Fischer [1, Theorem 6.2] will play a key role later on:

Lemma 1 (Csaki and Fischer): Suppose A_n and B_n , $n \in \mathbb{N}$, are σ -fields, and the σ -fields $A_n \vee B_n$, $n \in \mathbb{N}$ are independent. Then

$$\rho\left(\bigvee_{n \in \mathbb{N}} A_n, \bigvee_{n \in \mathbb{N}} B_n\right) = \sup_{n \in \mathbb{N}} \rho(A_n, B_n).$$

For a proof see Witsenhausen [7, Theorem 1]. (In that context there were only finitely many pairs of σ -fields, but the extension to Lemma 1 is elementary.)

Next, for any family Z of random variables let $B(Z)$ denote the σ -field of events generated by Z .

Lemma 2: Suppose c_1, c_2, c_3, \dots is a sequence of positive numbers. Then there exists a non-degenerate sequence $(Y_k, k \in \mathbb{Z})$ of real-valued random variables such that the following two statements hold:

- (i) $\forall n \in \mathbb{N}, \forall K \in \mathbb{Z}, \rho(B(Y_k, k \leq K), B(Y_k, k \geq K+n)) \leq c_n.$
- (ii) $\forall K \in \mathbb{Z}, Y_K$ is measurable with respect to $\bigvee_{n=1}^{\infty} (B(Y_k, k \leq -n) \vee B(Y_k, k \geq n)).$

The sequence (Y_k) that we shall construct for Lemma 2 will not be stationary, but it will obviously have the other properties needed for Theorem 1. In Section 3 the stationary sequence (X_k) for Theorem 1 will be constructed with the aid of a countable family of independent copies of the sequence (Y_k) here. The rest of Section 2 is devoted to the proof of Lemma 2.

Proof of Lemma 2: Without loss of generality we assume that

$$1 \geq c_1 \geq c_2 \geq c_3 \geq \dots > 0 \quad (2.1)$$

For each $n \in \mathbb{N}$ define

$$r_n = c_{2n} \quad (2.2)$$

(The purpose of (2.2) will become clear later on.)

On some probability space, let $(V_{k,m}: k \in \mathbb{Z}, m \in \mathbb{N})$ be an array of random variables with the following three properties:

(2.3) For each (k,m) ,

$$P(V_{k,m} = -1) = P(V_{k,m} = 1) = 1/2.$$

(2.4) The random variables $V_{0,m}$, $m \in \mathbb{N}$ and the random vectors $(V_{-n,m}, V_{n,m})$, $n \in \mathbb{N}$, $m \in \mathbb{N}$ are all independent of each other.

(2.5) For each $n \in \mathbb{N}$, $m \in \mathbb{N}$, $i \in \{-1,1\}$, $j \in \{-1,1\}$,

$$P(V_{-n,m}=i, V_{n,m}=j) = (1 + ijr_n)/4.$$

Let $h: \mathbb{N} \rightarrow \mathbb{N}$ be a function such that for each $\ell \in \mathbb{N}$ the following two statements hold:

(2.6) There exist infinitely many odd integers $m \in \mathbb{N}$ such that $h(m) = \ell$.

(2.7) There exist infinitely many even integers $m \in \mathbb{N}$ such that $h(m) = \ell$.

For each $\ell \in \mathbb{N}$, $J \in \mathbb{N}$, define the two sets $S(\ell, J, \text{odd})$ and $S(\ell, J, \text{even})$ as follows: $S(\ell, J, \text{odd})$ (resp. $S(\ell, J, \text{even})$) is the set of the least J positive odd (resp. even) integers m such that $h(m) = \ell$.

By (2.5) we have that for each $n \in \mathbb{N}$, $m \in \mathbb{N}$, $EV_{-n,m}V_{n,m} = r_n$; and by (2.4) and

the strong law of large numbers, for each $n \in \mathbb{N}$, $\ell \in \mathbb{N}$,

$$\lim_{J \rightarrow \infty} (1/J) \cdot \sum_{m \in S(\ell, J, \text{odd})} V_{-n, m} V_{n, m} = r_n \text{ a.s. and}$$

$$\lim_{J \rightarrow \infty} (1/J) \cdot \sum_{m \in S(\ell, J, \text{even})} V_{-n, m} V_{n, m} = r_n \text{ a.s.}$$

Deleting if necessary a null-set from the probability space on which the array $(V_{k, m})$ is defined, we assume that for every sample point ω in this probability space the following two statements hold:

$$(2.8) \quad \text{For every } k \in \mathbb{Z}, m \in \mathbb{N}, V_{k, m}(\omega) \in \{-1, 1\},$$

$$(2.9) \quad \text{For every } n \in \mathbb{N}, \ell \in \mathbb{N},$$

$$\lim_{J \rightarrow \infty} (1/J) \cdot \sum_{m \in S(\ell, J, \text{odd})} V_{-n, m}(\omega) \cdot V_{n, m}(\omega) = r_n \text{ and}$$

$$\lim_{J \rightarrow \infty} (1/J) \cdot \sum_{m \in S(\ell, J, \text{even})} V_{-n, m}(\omega) \cdot V_{n, m}(\omega) = r_n.$$

Let us digress for a moment to explain what will occur. Based on the array $(V_{k, m})$, another array $(W_{k, m})$ of $\{-1, 1\}$ -valued random variables will be defined, and for each $k \in \mathbb{Z}$ the r.v. Y_k in Lemma 2 will simply be a one-to-one bimeasurable function of $(W_{k, 1}, W_{k, 2}, W_{k, 3}, \dots)$. For each $n \in \mathbb{N}$, $\ell \in \mathbb{N}$, the r.v. $W_{-n, \ell}$ will be a function of the r.v.'s $(W_{-n-1, m}, W_{n+1, m}; m \text{ odd}, h(m) = \ell)$, and $W_{n, \ell}$ will be a function of $(W_{-n-1, m}, W_{n+1, m}; m \text{ even}, h(m) = \ell)$. In the end this will easily imply property (ii) in Lemma 2, but it has to be set up so that in the end property (i) of Lemma 2 will also hold.

Now let us define the array $(W_{k, m}; k \in \mathbb{Z}, m \in \mathbb{N})$. First,

$$\forall k \leq 0, \forall m \in \mathbb{N}, W_{k, m} := V_{k, m} \quad (2.10)$$

The definition of the other random variables $W_{k, m}$ will be recursive. For each

$n \geq 0$, once $(W_{k,m} : k \leq n, m \in \mathbb{N})$ is defined, define $W_{n+1,m}, m \in \mathbb{N}$ as follows:

$$\begin{aligned} \forall \text{ odd } m \in \mathbb{N}, \quad W_{n+1,m} &:= W_{-n,h(m)} \cdot V_{n+1,m} \\ \forall \text{ even } m \in \mathbb{N}, \quad W_{n+1,m} &:= W_{n,h(m)} \cdot V_{n+1,m} \end{aligned} \quad (2.11)$$

This completes the recursive definition.

In order to make our notation more compact, let us define for each $k \in \mathbb{Z}$ the random sequences V_k and W_k by $V_k := (V_{k,1}, V_{k,2}, V_{k,3}, \dots)$ and $W_k := (W_{k,1}, W_{k,2}, W_{k,3}, \dots)$.

By (2.8), (2.10), (2.11), and an induction argument, we have the following two facts:

(2.12) For every $k \in \mathbb{Z}, m \in \mathbb{N}$, the r.v. $W_{k,m}$ takes its values in $\{-1,1\}$ (at every sample point in our probability space).

(2.13) For every $n = 0,1,2,\dots$ the random sequences W_{-n} and W_n are measurable functions of $(V_{-n}, V_{-n+1}, \dots, V_n)$.

Claim 1: For each $n \geq 0$ the random sequences W_{-n} and W_n are each a measurable function of (W_{-n-1}, W_{n+1}) .

Proof of Claim 1: Let $n \geq 0$ and $\ell \in \mathbb{N}$ be arbitrary but fixed. Then at every sample point in our probability space, by (2.10), (2.11), the definition of $S(\ell, J, \text{odd})$, and (2.9),

$$\begin{aligned} & \lim_{J \rightarrow \infty} (1/J) \cdot \sum_{m \in S(\ell, J, \text{odd})} W_{-n-1,m} W_{n+1,m} = \\ &= \lim_{J \rightarrow \infty} (1/J) \cdot \sum_{m \in S(\ell, J, \text{odd})} V_{-n-1,m} \cdot W_{-n,h(m)} V_{n+1,m} = \\ &= W_{-n,\ell} \cdot \lim_{J \rightarrow \infty} (1/J) \cdot \sum_{m \in S(\ell, J, \text{odd})} V_{-n-1,m} V_{n+1,m} = W_{-n,\ell} \cdot r_{n+1}. \end{aligned}$$

Hence $W_{-n,\ell}$ is a measurable function of (W_{-n-1}, W_{n+1}) . The same is true for

$W_{n,\ell}$ by a similar argument using $S(\ell, J, \text{even})$. Claim 1 follows.

By Claim 1 and an induction argument we have the following:

Claim 2: For each $K \in \mathbb{Z}$, the random sequence W_K is measurable with respect to $\bigcap_{n=1}^{\infty} (B(W_K, k \leq -n) \vee B(W_K, k \geq n))$.

This will give us (ii) of Lemma 2 later on. Now we have to do some work to prepare for (i) of Lemma 2.

Claim 3: Suppose $n \in \mathbb{N}$, $m \in \mathbb{N}$, and Z is a random variable independent of $(V_{-n,m}, V_{n,m})$ and taking its values in $\{-1, 1\}$. Then the following two statements hold:

- (i) The joint distribution of $(Z \cdot V_{-n,m}, Z \cdot V_{n,m})$ is the same as that of $(V_{-n,m}, V_{n,m})$.
- (ii) $\rho(B(Z \cdot V_{-n,m}), B(Z \cdot V_{n,m})) = \rho(B(V_{-n,m}), B(V_{n,m})) = r_n$.

Proof of Claim 3: Part (i) follows from (2.5) and an elementary calculation. The first equality in (ii) follows from (i); we only need to verify the second equality in (ii).

Suppose f (resp. g) is a non-degenerate function of $V_{-n,m}$ (resp. $V_{n,m}$). Since $V_{-n,m}$ takes only two values $\{-1$ and $\}$

Claim 4: Suppose $n \geq 0$, and Z_1, Z_2, Z_3, \dots are $B(W_k, -n \leq k \leq n)$ -measurable random variables taking their values in $\{-1, 1\}$. Then the σ -fields $B(W_k, -n \leq k \leq n)$ and $B(Z_J \cdot V_{-n-1,J}, Z_J \cdot V_{n+1,J})$, $J = 1, 2, 3, \dots$ are all

independent of each other.

Proof of Claim 4: Let $J \geq 1$ be arbitrary but fixed. Define the σ -field

$$A := B(W_k, -n \leq k \leq n) \vee \left(\bigvee_{h=1}^{J-1} B(Z_h \cdot V_{-n-1,h}, Z_h \cdot V_{n+1,h}) \right). \quad (\text{In the case } J=1$$

define $A := B(W_k, -n \leq k \leq n)$.) To prove Claim 4 it suffices to prove that the random vector $(Z_J \cdot V_{-n-1,J}, Z_J \cdot V_{n+1,J})$ is independent of A .

Let $A \in \mathcal{A}$ be an arbitrary fixed event such that $P(A) > 0$. Let i, j each be an arbitrary fixed element of $\{-1, 1\}$. It suffices to prove that

$$\begin{aligned} P(Z_J \cdot V_{-n-1,J} = i, Z_J \cdot V_{n+1,J} = j | A) &= \\ &= P(Z_J \cdot V_{-n-1,J} = i, Z_J \cdot V_{n+1,J} = j). \end{aligned} \quad (2.14)$$

By (2.13) the random sequences $W_{-n}, W_{-n+1}, \dots, W_n$ are (measurable) functions of $V_{-n}, V_{-n+1}, \dots, V_n$. Consequently A is an element of, and Z_J is measurable with respect to, the σ -field

$B(V_k, -n \leq k \leq n) \vee B(V_{-n-1,h}, V_{n+1,h}; 1 \leq h \leq J-1)$; and $(V_{-n-1,J}, V_{n+1,J})$ is independent of this σ -field by (2.4). Consequently, for each $z \in \{-1, 1\}$ such that $P(A \cap \{Z_J = z\}) > 0$, one has that

$$\begin{aligned} P(Z_J \cdot V_{-n-1,J} = i, Z_J \cdot V_{n+1,J} = j | A \cap \{Z_J = z\}) &= \\ &= P(V_{-n-1,J} = i/z, V_{n+1,J} = j/z | A \cap \{Z_J = z\}) \\ &= P(V_{-n-1,J} = i/z, V_{n+1,J} = j/z) \\ &= (1 + r_{n+1} ij/z^2) = (1 + r_{n+1} ij) \\ &= P(Z_J \cdot V_{-n-1,J} = i, Z_J \cdot V_{n+1,J} = j). \end{aligned}$$

(We are using (2.5) and Claim 3(i).) Equation (2.14) now follows by a simple

argument, and this completes the proof of Claim 4.

In applying Claim 4 in what follows, we shall sometimes implicitly take $Z_j := 1$, in which case $B(Z_j V_{-n-1,j}, Z_j V_{n+1,j}) = B(V_{-n-1,j}, V_{n+1,j})$.

Claim 5: For each $n \in \mathbb{N}$ the following two statements hold:

- (i) $\rho(B(W_k, -n \leq k \leq n-1), B(W_n)) \leq r_n$.
- (ii) $\rho(B(W_{-n}), B(W_k, -n+1 \leq k \leq n)) \leq r_n$.

Proof of Claim 5: Let $n \in \mathbb{N}$ be arbitrary but fixed.

Proof of Claim 5(i): Define the σ -fields A_j and B_j , $j = 0, 1, 2, \dots$ as follows: $A_0 := B(W_k, -n+1 \leq k \leq n-1)$ and $B_0 := \{\Omega, \phi\}$. For odd $j \geq 1$, $A_j := B(W_{-n+1, h(j)} \cdot V_{-n, j})$ and $B_j := B(W_{-n+1, h(j)} \cdot V_{n, j})$. For even $j \geq 2$, $A_j := B(W_{n-1, h(j)} \cdot V_{-n, j})$ and $B_j := B(W_{n-1, h(j)} \cdot V_{n, j})$. (Here of course Ω denotes the sample space.) Define the σ -fields A and B by

$$A := \bigvee_{j=0}^{\infty} A_j \quad \text{and} \quad B := \bigvee_{j=0}^{\infty} B_j$$

For each odd $j \in \mathbb{N}$, by (2.10) and (2.12),

$W_{-n, j} = V_{-n, j} = (W_{-n+1, h(j)} \cdot V_{-n, j}) \cdot W_{-n+1, h(j)}$, and since $(W_{-n+1, h(j)} \cdot V_{-n, j})$ and $W_{-n+1, h(j)}$ are each A -measurable, $W_{-n, j}$ is A -measurable. Similarly, for each even $j \in \mathbb{N}$, $W_{-n, j}$ is A -measurable. Hence W_{-n} is A -measurable. (Of course W_{-n+1}, \dots, W_{n-1} are also A -measurable.)

Also, for each $j \geq 1$, $B_j = B(W_{n, j})$ by (2.11); and hence $B = B(W_n)$. Hence

$$\rho(B(W_{-n}, W_{-n+1}, \dots, W_{n-1}), B(W_n)) \leq \rho(A, B) = \sup_{j \geq 0} \rho(A_j, B_j) = r_n.$$

Here the inequality is trivial, the first equality holds by Claim 4 and Lemma 1, and the second equality holds by Claim 3. This completes the proof of (i).

Proof of Claim 5(ii): This is analogous to the proof of (i), but with a few changes. One defines the σ -fields A_j and B_j as follows:

$$A_0 := (\Omega, \phi) \quad \text{and} \quad B_0 := B(W_{-n+1}, \dots, W_{n-1})$$

$$\text{For all } j \geq 1, \quad A_j := B(V_{-n,j}) \quad \text{and} \quad B_j := B(V_{n,j})$$

Again one defines $A := \bigvee_{j=0}^{\infty} A_j$ and $B := \bigvee_{j=0}^{\infty} B_j$. Then one observes that $A := B(W_{-n})$

and that W_n (as well as W_{-n+1}, \dots, W_{n-1}) is B -measurable. The proof of (ii) is then completed in the same way as (i).

Claim 6: For each $n \in \mathbb{N}$ the following two statements hold:

$$(i) \quad \rho(B(W_k, k \leq n-1), B(W_k, k \geq n)) \leq r_n.$$

$$(ii) \quad \rho(B(W_k, k \leq -n), B(W_k, k \geq -n+1)) \leq r_n.$$

Proof of Claim 6: Let $n \in \mathbb{N}$ be arbitrary but fixed.

Proof of Claim 6(i): It suffices to prove that $\forall N \geq n$,

$\rho(B(W_k, -N \leq k \leq n-1), B(W_k, n \leq k \leq N)) \leq r_n$. We shall show this by induction on N . By Claim 5, $\rho(B(W_k, -n \leq k \leq n-1), B(W_n)) \leq r_n$. Now we only need to carry out the induction step.

Assume that $N \geq n$ is such that

$$\rho(B(W_k, -N \leq k \leq n-1), B(W_k, n \leq k \leq N)) \leq r_n. \quad \text{To prove Claim 6(i) it suffices}$$

to show that $\rho(B(W_k, -N-1 \leq k \leq n-1), B(W_k, n \leq k \leq N+1)) \leq r_n$. The proof will be somewhat similar to that of Claim 5.

Define the σ -fields A_j and B_j , $j=0,1,2,\dots$ as follows:

$$A_0 := B(W_k, -N \leq k \leq n-1) \quad \text{and} \quad B_0 := B(W_k, n \leq k \leq N)$$

For each odd $j \geq 1$, $A_j := B(W_{-N,h(j)} \cdot V_{-N-1,j})$ and

$$B_j := B(W_{-N,h(j)} \cdot V_{N+1,j})$$

For each even $j \geq 2$, $A_j := B(V_{-N-1,j})$ and $B_j := B(V_{N+1,j})$

Define the σ -fields A and B by

$$A := \bigvee_{j=0}^{\infty} A_j \quad \text{and} \quad B := \bigvee_{j=0}^{\infty} B_j$$

For each odd $j \in \mathbb{N}$, by (2.10) and (2.12),

$W_{-N-1,j} = V_{-N-1,j} = (W_{-N,h(j)} \cdot V_{-N-1,j}) \cdot W_{-N,h(j)}$, which is a product of two

A -measurable r.v.'s, and hence $W_{-N-1,j}$ is A -measurable. For each even $j \in \mathbb{N}$,

$W_{-N-1,j} = V_{-N-1,j}$, which is A -measurable. Hence W_{-N-1} is A -measurable.

For each odd $j \in \mathbb{N}$, by (2.11), $W_{N+1,j} = W_{-N,h(j)} \cdot V_{N+1,j}$, which is B -measurable. For each even $j \in \mathbb{N}$, $W_{N+1,j} = W_{N,h(j)} \cdot V_{N+1,j}$, which is a product of B -measurable functions and is therefore B -measurable. Hence W_{N+1} is B -measurable.

Consequently,

$$\begin{aligned} \rho(B(W_k, -N-1 \leq k \leq n-1), B(W_k, n \leq k \leq N+1)) &\leq \rho(A, B) = \sup_{j \geq 0} \rho(A_j, B_j) \leq \\ &\leq \max \{r_n, r_{N+1}\} = r_n. \end{aligned}$$

Here the first inequality is trivial, we are using Claim 4 and Lemma 1 to obtain the first equality, and then Claim 3 and the induction hypothesis to obtain the second inequality, and finally (2.1) and (2.2) to obtain the second equality. This completes the induction step, and Claim 6(i) is proved.

Proof of Claim 6(ii): This is essentially identical to the proof of (i). One has to change the definition of A_0 and B_0 to $A_0 := B(W_k, -N \leq k \leq -n)$ and $B_0 := B(W_k, -n+1 \leq k \leq N)$.

Claim 7: If $n \in \mathbb{N}$ and $K \in \mathbb{Z}$ then $\rho(B(W_k, k \leq K), B(W_k, k \geq K+n)) \leq c_n$.

Proof of Claim 7: Let us first consider the case where n is odd, say $n = 2m-1$ where $m \in \mathbb{N}$. Then either $K \leq -m$ or $K+n \geq m$. If $K \leq -m$ then $-2K \geq 2m > n$, and

$$\begin{aligned} & \rho(B(W_k, k \leq K), B(W_k, k \geq K+n)) \leq \\ & \leq \rho(B(W_k, k \leq K), B(W_k, k \geq K+1)) \\ & \leq r_{-K} = c_{-2K} \leq c_n \end{aligned}$$

by Claim 6(ii), (2.2), and (2.1). If instead $K+n \geq m$ then a similar argument (using Claim 6(i)) works. The proof for even n is similar to that for odd n .

The rest of the proof of Lemma 2 is now a trivial, cosmetic formality. Let $f: \{-1,1\} \times \{-1,1\} \times \{-1,1\} \times \dots \rightarrow \mathbb{R}$ be a bimeasurable isomorphism. (It is well known that such an f exists.) Define the sequence $(Y_k, k \in \mathbb{Z})$ by $Y_k := f(W_k) \forall k \in \mathbb{Z}$. Properties (i) and (ii) of Lemma 2 now follow immediately from Claims 7 and 2 respectively. This completes the proof of Lemma 2.

3. Proof of Theorem 1

Let c_1, c_2, c_3, \dots be as in the statement of Theorem 1. Our first step is to construct countably many copies of the probability space and random sequence $(Y_k, k \in \mathbb{Z})$ from Lemma 2. Then on the product probability space we obtain an array $(Y_k^{(j)}: k \in \mathbb{Z}, j \in \mathbb{Z})$ of real-valued random variables with the following three properties:

(3.1) The sequences $(Y_k^{(j)}, k \in \mathbb{Z}), j = \dots, -1, 0, 1, \dots$, are non-degenerate and are independent of each other, and they have the same distribution on $\dots \times \mathbb{R} \times \mathbb{R} \times \mathbb{R} \times \dots$.

(3.2) For each $j \in \mathbb{Z}, n \in \mathbb{N}, K \in \mathbb{Z}, \rho(B(Y_k^{(j)}, k \leq K), B(Y_k^{(j)}, k \geq K+n)) \leq c_n$.

(3.3) For each $j \in \mathbb{Z}, K \in \mathbb{Z}$, the r.v. $Y_K^{(j)}$ is measurable with respect to $\bigcap_{n=1}^{\infty} (B(Y_k^{(j)}, k \leq -n) \vee B(Y_k^{(j)}, k \geq n))$.

Let $g: \dots \times \mathbb{R} \times \mathbb{R} \times \mathbb{R} \times \dots \rightarrow \mathbb{R}$ be a bimeasurable isomorphism. (It is well known that such a function g exists.) Define the sequence $(X_k, k \in \mathbb{Z})$ of real-valued random variables as follows:

$$\forall k \in \mathbb{Z}, X_k := g(\dots, Y_{-1}^{(k-1)}, Y_0^{(k)}, Y_1^{(k+1)}, \dots) \quad (3.4)$$

By a simple argument using (3.1), this sequence (X_k) is non-degenerate and strictly stationary.

For each $n \in \mathbb{N}$, by (3.4),

$$\begin{aligned} B(X_k, k \leq -n) \vee B(X_k, k \geq n) &= \\ &= \bigvee_{j \in \mathbb{Z}} (B(Y_k^{(j)}, k \geq j+n) \vee B(Y_k^{(j)}, k \leq j-n)) \end{aligned}$$

Hence for each $n \in \mathbb{N}$, $K \in \mathbb{Z}$, the r.v. X_K is measurable with respect to $B(X_k, k \leq -n) \vee B(X_k, k \geq n)$, by a simple argument using (3.3) and (3.4). Hence the sequence (X_k) is bilaterally deterministic.

For each $n \in \mathbb{N}$,

$$\begin{aligned} \rho(n) &:= \rho(B(X_k, k \leq 0), B(X_k, k \geq n)) \\ &= \rho(\vee_{j \in \mathbb{Z}} B(Y_k^{(j)}, k \geq j), \vee_{j \in \mathbb{Z}} B(Y_k^{(j)}, k \leq j-n)) \\ &= \sup_{j \in \mathbb{Z}} \rho(B(Y_k^{(j)}, k \geq j), B(Y_k^{(j)}, k \leq j-n)) \leq c_n \end{aligned}$$

by (3.4), (3.1), Lemma 1, and (3.2). This completes the proof of Theorem 1.

Acknowledgments: The author is indebted to Henry Berbee for suggesting this topic, and for calling attention to the earlier references. For their hospitality during which parts of this work were completed, the author also thanks Włodzimierz and Grazyna Bryc as well as the Mathematical Centre in Amsterdam and the Center for Stochastic Processes in Chapel Hill.

REFERENCES

1. P. Csaki and J. Fischer. On the general notion of maximal correlation. Magyar Tud. Akad. Mat. Kutato Int. Kozl. 8 (1963) 27-51.
2. V.M. Gurevič. On one-sided and two-sided regularity of stationary random processes. Soviet Math. Dokl. 14 (1973) 804-808.
3. I.A. Ibragimov. A note on the central limit theorem for dependent random variables. Theory Probab. Appl. 20 (1975) 135-141.
4. I.A. Ibragimov and Y.V. Linnik. Independent and Stationary Sequences of Random Variables. Wolters-Noordhoff, Groningen, 1971.
5. R.A. Olshen. The coincidence of measure algebras under an exchangeable probability. Z. Wahrsch. verw. Gebiete 18 (1971) 153-158.
6. D.S. Ornstein and B. Weiss. Every transformation is bilaterally deterministic. Israel J. Math. 21 (1975) 154-158.
7. H.S. Witsenhausen. On sequences of pairs of dependent random variables. SIAM J. Appl. Math. 28 (1975) 100-113.

END

FILMED

5-85

DTIC